

Understanding Society User Support - Support #1733

Covid Survey - Measuring Internal Migration

07/25/2022 02:52 PM - Morgan Ward

Status:	Resolved	Start date:	07/25/2022
Priority:	Normal	% Done:	100%
Assignee:			
Category:	Data management		
Description			
Hello,			
I am attempting to measure 3-monthly internal migration moves between UK regions during the Covid pandemic, using your Covid survey data.			
To do this, I have created a variable to identify where movements have taken place between waves, e.g., for moves between May - July 2020: gen regi_moves = 1 if cd_gor_dv != cb_gor_dv			
However, when I longitudinally weight these regional moves using betaindin_lw, the unweighted counts of moves are significantly reduced. E.g., the unweighted count of 147 moves into the South East between May - July 2020, is reduced to a weighted count of 3.324 moves.			
Is there a way of producing valid weights for the analysis of moves between Waves please, which do not have such reduced counts? I was hoping to compare sequential Waves, excluding Waves 1 and 3. So comparing Waves 2 - 4, 4 - 5, 5 - 6 etc. The reduction in counts prevents me from generalizing rates of internal migration moves with any accuracy.			
Any help will be greatly appreciated.			
Thank you			

History

#1 - 07/25/2022 03:45 PM - Understanding Society User Support Team

- File regi_moves.jpg added
- Category set to Data management
- Status changed from New to Feedback
- % Done changed from 0 to 80
- Private changed from Yes to No

Dear Morgan,

I suspect that you have incorrectly calculated the number of moves. I managed to get 147 moves to South East between May and July when I ran your syntax creating regi_moves on a datafile that includes all respondents after merging cb_indresp and cd_indresp, that is, including the respondents who participated in wave 2 only (so did not participate in wave 4) and the respondents who participated in wave 4 only (so not in wave 2). For such cases regi_moves equals 1 but that's a Stata artefact (see the screenshot), not a correctly identified move. The correct total number of moves between these time points is 16 (that is, calculated on a sample where all respondents participated in both waves). As longitudinal weights are calculated only for people who continuously participated in the study, that is, in all waves you are interested in (in this case w2, 3, and 4), when you weighted using cd_betaindin_lw Stata automatically excluded all these artificial moves from the calculation (as cd_betaindin_lw for these cases is either missing or equals 0).

I hope this helps.

Best wishes,
Piotr
UKHLS User Support Team

#2 - 07/25/2022 03:49 PM - Morgan Ward

Dear Piotr,

This was very helpful, thank you.

Best wishes,

Morgan

#3 - 10/20/2022 08:32 AM - Understanding Society User Support Team

- Status changed from *Feedback* to *Resolved*

- % Done changed from 80 to 100

Files

regi_moves.jpg	25.2 KB	07/25/2022	Understanding Society User Support Team
----------------	---------	------------	---