

## Understanding Society User Support - Support #349

### GLYCATED HAEMOGLOBIN (HbA1c)

03/16/2015 05:37 PM - Rasha Alfawaz

<b>Status:</b>	Closed	<b>Start date:</b>	03/16/2015
<b>Priority:</b>	Urgent	<b>% Done:</b>	100%
<b>Assignee:</b>			
<b>Category:</b>	Derived variables		
<b>Description</b> Hi, I am interested in analysing HbA1c data which is available at xlabblood_ns dataset. In the Biomarker User Guide it was mentioned that a number of factors are associated with decreased HbA1c measurements <sup>19</sup> - chronic liver disease, taking aspirin (BNF chapter 2.9; derived variable bnf7_aspirin) and anti-inflammatory medications (BNF chapter 10.1; derived variable: bnf7_antiinflam). High levels of triglycerides may lead to artefactually low measurements. However those derived variables are not available (aspirin and anti-inflammatory) I looked in the (indresp_ns) file and nothing was there too. My query is what are the variables of those medications and where can I find them as I need to adjust for in the model am using.  Best regards			

#### History

##### #1 - 03/17/2015 09:57 AM - Redmine Admin

- Status changed from New to In Progress
- % Done changed from 0 to 50

can be found on xindresp\_ns - documented around here:

[https://www.understandingsociety.ac.uk/documentation/health-assessment/dataset-documentation/wave/xwave/datafile/xindresp\\_ns/variable/bnf7\\_aspirin](https://www.understandingsociety.ac.uk/documentation/health-assessment/dataset-documentation/wave/xwave/datafile/xindresp_ns/variable/bnf7_aspirin)

On behalf of the team,  
Jakob

##### #2 - 03/26/2015 04:22 PM - Redmine Admin

- Status changed from In Progress to Closed
- % Done changed from 50 to 100