

Understanding Society User Support - Support #919

weights for pooled cross-sections over waves (a)-(g)

02/14/2018 11:15 AM - Nico Ochmann

Status:	Resolved	Start date:	02/14/2018
Priority:	High	% Done:	100%
Assignee:	Nico Ochmann		
Category:	Weights		
Description Dear Peter, I have a quick follow-up question with regard to my past issue #839 . You wrote the following below: A simpler approach would be to select based on current wave GOR. This will not remove the sample imbalance problem, but should reduce it to a minimum, so may be good enough. For me to implement this simpler approach and to minimize the sample imbalance, you suggest to drop Scot, Wales, NI for the current wave g only? Or do you mean to drop all Scot, Wales, NI for all seven waves? I would appreciate a short reply and sorry about the very delayed follow-up question. Best. Nico			

History

#1 - 02/14/2018 11:42 AM - Peter Lynn

- Assignee changed from Peter Lynn to Nico Ochmann
- % Done changed from 0 to 50

Nico,

By "current wave" I meant wave 1 in the case of the wave 1 sample, wave 2 for the wave 2 sample, etc. So, for each of the 6 waves you are pooling, select based on GOR for that wave. That way your analysis will represent - for each year - people living in England at that time.

Peter

#2 - 02/14/2018 11:42 AM - Peter Lynn

- Status changed from New to Feedback
- Private changed from Yes to No

#3 - 04/05/2018 12:35 PM - Stephanie Auty

- Status changed from Feedback to Resolved
- % Done changed from 50 to 100

From: Nico Ochmann [mailto:nico.ochmann@manchester.ac.uk]

Sent: 14 February 2018 12:24

To: usersupport@understandingsociety.ac.uk

Subject: RE: [Understanding Society User Support - Support [#919](#)] weights for pooled cross-sections over waves (a)-(g)

Hi Peter,

I appreciate your prompt reply very much. It is very clear to me now.

Apologies for the confusion on my part.

Have a great day.

Nico